

Effect of storage conditions on the assessment of bacterial community structure in soil and human-associated samples

Christian L. Lauber¹, Nicholas Zhou², Jeffrey I. Gordon³, Rob Knight⁴ & Noah Fierer^{1,2}

¹Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA; ²Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA; ³Center for Genome Sciences, Washington University School of Medicine, St. Louis, MO, USA; and ⁴Department of Chemistry and Biochemistry, University of Colorado, Boulder, CO, USA

Correspondence: Christian L. Lauber, Cooperative Institute for Research in Environmental Sciences, University of Colorado, UCB 216, Boulder, CO 80309-0216, USA. Tel.: +1 303 492 2099; fax: +1 303 492 1149; e-mail: chris.lauber@gmail.com

Received 27 October 2009; revised 2 March 2010; accepted 12 March 2010.
Final version published online April 2010.

DOI:10.1111/j.1574-6968.2010.01965.x

Editor: Elizabeth Baggs

Keywords

microbial community storage conditions; environmental and human metagenomic studies; barcoded bacterial 16S rRNA gene pyrosequencing; phylogenetic- and taxonomic-based community analyses; soil; human fecal and human skin microbiota.

Introduction

The treatment and handling of samples after collection is a critical aspect of a study design when using DNA-based methods to compare the composition and diversity of microbial communities from environmental samples. It is widely assumed that microbial DNA must be extracted from the samples immediately after collection or, if this is not possible, that samples must be frozen (Rochelle *et al.*, 1994). Samples stored at room temperature even for a short period before DNA is extracted are often considered unfit for downstream analyses because of changes to the microbial community. Although these assumptions are widespread, few and conflicting studies have directly tested the influence of storage conditions on DNA-based bacterial community analyses. For example, Dolfig *et al.* (2004) and Klammer *et al.* (2005) used DNA fingerprinting methods to show that the overall structure of soil bacterial communities was not strongly affected by

Abstract

Storage conditions are considered to be a critical component of DNA-based microbial community analysis methods. However, whether differences in short-term sample storage conditions impact the assessment of bacterial community composition and diversity requires systematic and quantitative assessment. Therefore, we used barcoded pyrosequencing of bacterial 16S rRNA genes to survey communities, harvested from a variety of habitats [soil, human gut (feces) and human skin] and subsequently stored at 20, 4, –20 and –80 °C for 3 and 14 days. Our results indicate that the phylogenetic structure and diversity of communities in individual samples were not significantly influenced by the storage temperature or the duration of storage. Likewise, the relative abundances of most taxa were largely unaffected by temperature even after 14 days of storage. Our results indicate that environmental factors and biases in molecular techniques likely confer greater amounts of variation to microbial communities than do differences in short-term storage conditions, including storage for up to 2 weeks at room temperature. These results suggest that many samples collected and stored under field conditions without refrigeration may be useful for microbial community analyses.

storage conditions. Likewise, Roesch *et al.* (2009) reported only modest shifts in the bacterial diversity in only one of four human gut samples after 72 h of storage at room temperature. In contrast, both Tzeneva *et al.* (2009) and Ott *et al.* (2004) observed significant effects of storage conditions on the composition and diversity of microbial communities in soil and human gut samples, respectively. Nechvatal *et al.* (2008) tested a series of room-temperature methods for preserving stool for up to 5 days and found considerable variability among methods, but tested each sample with a different stool specimen, limiting their ability to draw general conclusions because of the large degree of variability among stool specimens from different human subjects.

Because of the diversity of the samples, conditions tested and analytic methods used, we still lack a comprehensive understanding of how and whether storage of samples before DNA extraction impacts bacterial community analyses and the magnitude of these potential storage effects. In

particular, we do not know whether variation in storage conditions (temperature and length of storage) influences our ability to resolve differences in the bacterial community composition and diversity between samples. To address these knowledge gaps, we analyzed bacterial communities in soil, human skin and human fecal samples stored for different amounts of time and at varying temperatures using a barcoded pyrosequencing procedure, with the sequence data from each sample analyzed using both phylogenetic and taxonomic community analysis procedures.

Methods

Sample collection and storage conditions

Microbial communities were sampled from three distinct habitats: surface soils, human skin and human feces. Fecal samples (Fecal 1 and 2; *c.* 100 g each) were donated by two anonymous male participants. Immediately after collection, each sample was homogenized by stirring with a sterile spatula without added buffer in a sterile container. Replicate subsamples ($n=24$) of each homogenized fecal sample were obtained by inserting sterile cotton swabs into each sample, and then placing the swab into its own separate dry, sterile 15-mL conical tube. Soil was collected ($3\text{--}2.5 \times 10$ cm cores) from two locations on the campus of the University of Colorado ($40^\circ 0'N$, $105^\circ 16'W$) in June 2009. One set of cores was collected from underneath a *Pinus ponderosa* tree (Soil 1), while the other was collected from an irrigated lawn (Soil 2). Replicate cores were composited and sieved through a 2-mm mesh and thoroughly homogenized by hand. From these two soil samples, forty-eight 1-g subsamples ($n=24$ per sample) were each placed in 1.5-mL centrifuge tubes. Skin samples were taken from the axillae (armpits) of one male and one female volunteer using sterile cotton swabs that had been premoistened in a sterile solution of 0.15 M NaCl and 0.01% Tween 20 (Paulino *et al.*, 2006; Fierer *et al.*, 2008). The axillary surface was swabbed for 30 s with all 24 swabs per individual at one time. The swabs were then placed in sterile 15-mL conical tubes for storage.

Replicate subsamples of each community type ($n=3$) were subsequently stored at 20, 4, -20 and -80°C for either 3 or 14 days before DNA extraction. All sample-treatment combinations (four storage temperatures; two storage times; six unique samples) were analyzed in triplicate as described in the next paragraph. Participants in the study gave informed consent under the sampling protocol approved by the University of Colorado Human Research Committee (protocol 1007.39).

DNA extraction, PCR, amplicon pooling and pyrosequencing

Genomic DNA was extracted from all samples using the MoBio Power Soil DNA extraction kit (MoBio, Carlsbad,

CA) as described previously (Fierer *et al.*, 2008; Lauber *et al.*, 2009). Note that samples were placed in bead tubes containing solution C1 and incubated at 65°C for 10 min, followed by 2 min of bead beating with the MoBio vortex adapter; the remaining steps of the extraction were performed as directed by the manufacturer. PCR amplification of bacterial 16S rRNA genes using primers directed at variable regions V1 and V2 (positions 27–338 according to the *Escherichia coli* 16S rRNA gene numbering scheme) was achieved following the protocol described in our earlier publications (Fierer *et al.*, 2008; Lauber *et al.*, 2009). Briefly, amplicons generated from three PCR reactions per sample were pooled to reduce per-PCR variability and purified using the MoBio Ultra Clean PCR cleanup kit according to the manufacturer's instructions and quantified (PicoGreen; Invitrogen, Carlsbad, CA). No-template PCR controls were also performed. PCR products generated from each subsample contained a sub-sample-specific, error-correcting barcode, which allowed us to assemble a single composite sample for pyrosequencing by combining equal amounts of amplicon from each subsample. The composite sample was then gel purified (Qiaquick gel Clean up kit, Qiagen, Valencia, CA) and precipitated with ethanol to remove any remaining contaminants. DNA was sequenced using a Roche 454 FLX pyrosequencer.

Sequence analysis

16S rRNA gene sequences were processed according to the methods described in our previous publications (Fierer *et al.*, 2008; Hamady *et al.*, 2008). Briefly, sequences < 200 or > 300 nt or with average quality scores of < 25 were removed from the dataset, as were those with uncorrectable barcodes, ambiguous bases, or if the bacterial 16S rRNA gene-specific primer was absent. Sequences were then assigned to the specific subsamples based on their unique 12 nt barcode and then grouped into phylotypes at the 97% level of sequence identity using cd-hit (Li & Godzik, 2006) with a minimum coverage of 97%. We chose to group the phylotypes at 97% identity because this matches the limits of resolution of pyrosequencing (Kunin *et al.*, 2010) and because the branch length so omitted contributes little to the tree and therefore to phylogenetic estimates of β diversity (Hamady *et al.*, 2009). A representative for each phylotype was chosen by selecting the most abundant sequence in the phylotype, with ties being broken by choosing the longest sequence. A phylogenetic tree of the representative sequences was constructed using the Kimura 2-parameter model in Fast Tree (Price *et al.*, 2009) after sequences were aligned with NAST (minimum 150 nt at 75% minimum identity) (DeSantis *et al.*, 2006a) against the GreenGenes database (DeSantis *et al.*, 2006b). Hypervariable regions were screened out of the alignment

using PH Lane mask (<http://greengenes.lbl.gov/>). Differences in the community composition for each pair of samples were determined from the phylogenetic tree using the weighted and unweighted UniFrac algorithms (Lozupone & Knight, 2005; Lozupone *et al.*, 2006). UniFrac is a tree-based metric that measures the distance between two communities as the fraction of branch length in a phylogenetic tree that is unique to one of the communities (as opposed to being shared by both). This method of community comparison accounts for the relative similarities and differences among phylotypes (or higher taxa) rather than treating all taxa at a given level of divergence as equal (Lozupone & Knight, 2008). Although UniFrac depends on a phylogenetic tree, it is relatively robust to differences in the tree reconstruction method or to the approximation of using phylotypes to represent groups of very similar sequences (Hamady *et al.*, 2009).

UniFrac calculates the unique fraction of branch length for a sample from a phylogenetic tree constructed from each pair of samples in a data set. Because the UniFrac metric is a phylogenetic estimate of community similarity, it avoids some of the problems associated with analyses that compare communities at arbitrarily defined levels of sequence similarity (Lozupone & Knight, 2008; Hamady & Knight, 2009). The phylogenetic diversity of each sample was determined from 1000 randomly selected sequences per sample using Faith's phylogenetic diversity metric (Faith's PD; Faith, 1992), which calculates the amount of branch length for each sample within the relaxed neighbor-joining tree. The taxonomic identity of each phylotype was determined using the RDP-II taxonomy (60% minimum threshold) (Cole *et al.*, 2005). All sequences have been deposited in the GenBank short read archive (accession number SRA012078.1).

Statistical analyses

The effect of temperature and length of storage on the relative taxon abundance (minimum 1% abundance per sample-treatment combination) was assessed using the Kruskal-Wallis test in SYSTAT 11.0 for sequences classified to the level of order (fecal and skin) or family (soil). Statistical differences in the overall community composition (UniFrac distances) were assessed within each sample type using the PERMANOVA package in PRIMER v6 using Sample, Day, Temperature and Day \times Temperature as the main factors. Pairwise UniFrac distances were visualized by non-metric multidimensional scaling in PRIMER v6 (Clarke & Warwick, 2001). Differences in Faith's PD due to the temperature and length of storage were assessed using the Kruskal-Wallis test.

Results and discussion

Sequence analysis and overall results

After eliminating low-quality sequences, the number of reads ranged from 1304 to 3022 per subsample, with an average of 2019 sequences per subsample and a total of 290 696 sequences for the data set. One subsample was excluded from the data set (Fecal 1 Day 14, 20 °C replicate 2) due to visible fungal growth before DNA extraction. Each sample type yielded a similar total number of bacterial 16S rRNA gene sequences (97 943 for feces, 97 527 for skin and 95 226 for soil). These distinct sample types harbored communities that were distinct with respect to their composition and diversity (Figs 1 and 2 and Tables 1 and 2). The human-derived samples tended to have few dominant phyla (three per sample type) accounting for 95–98% of the sequences: in contrast, in the soils, the six most abundant phyla accounted for only *c.* 83% of the sequences. Fecal samples were dominated by members of the phylum *Bacteroidetes* (62%) and the *Firmicutes* (35%), while skin samples had a relatively even distribution of *Firmicutes* (39%), *Bacteroidetes* (31%) and *Actinobacteria* (25%). Soil samples contained many phyla including the *Bacteroidetes* (32%), *Acidobacteria* (27%) and *Proteobacteria* [*Alphaproteobacteria* (10%), *Betaproteobacteria* (6.5%), *Gammmaproteobacteria* (5.2%) and *Deltaproteobacteria* (2.7%)]. The unique distribution of phyla was also seen in the overall community composition as NMDS visualization of pairwise UniFrac distances showed clustering by individual sample rather than temperature or length of storage (Fig. 1). Sample types also differed with respect to community diversity levels, with soil bacterial communities harboring the highest levels, followed by fecal and skin samples (Faith's PD = 40, 11 and 10, respectively). As noted below, each pair of subsamples within a given sample type had bacterial communities that differed with respect to their composition and diversity and these differences were irrespective of the storage conditions (see Table 1).

Storage effects: fecal samples

Bacterial communities in the fecal samples did not change appreciably with different storage conditions and retained their unique composition even after 14 days of storage (Fig. 1 and Table 1). Fecal 2 was dominated by the *Bacteroidaceae* (*c.* 75%), while Fecal 1 had a more even distribution of the six most abundant taxa across all temperatures (Fig. 2). Although the relative abundances of a few individual taxa were affected by temperature (Fig. 2), this did not have a significant effect on the overall community composition. The UniFrac distance between bacterial communities from the two hosts was significantly greater (PERMANOVA, $P = < 0.001$) for both weighted and unweighted UniFrac




Fig. 1. Nonmetric multidimensional scaling (NMDS) plots of UniFrac weighted and unweighted pairwise distances. The overall community composition was not affected by temperature or duration of storage for weighted UniFrac distances ($P > 0.1$ in all cases). Length of storage significantly affected the skin communities for the unweighted UniFrac metric ($P = 0.02$). The remaining unweighted UniFrac distances were not significantly different by day or temperature. Blue, sample 1; red, sample 2; open symbols, Day 3; closed symbols, Day 14; ▲, 20 °C; ■, 4 °C; ●, -20 °C; ◆, -80 °C.

than the distance between samples stored at different temperatures and durations ($P > 0.1$, Table 1). This minimal effect of storage on the overall structure of the communities is evident from Fig. 1, which shows that replicate samples tended to cluster by host. Likewise, the phylogenetic diversity of the fecal samples remained consistent across the temperatures (Table 2). Our results extend those reported by Roesch *et al.* (2009), who found minimal differences in community composition and relative taxon abundances after 72 h of storage at the one temperature tested (room temperature) for three of the four samples in their study.

In summary, even though we did observe shifts in the abundance of some taxa in our small sample set under different storage conditions, this did not mask interpersonal differences in the overall fecal bacterial community composition, and did not affect our ability to differentiate the host origin of the two fecal samples.

Skin samples

As with the fecal communities, each skin sample was distinctive in terms of dominant taxa and the overall community composition. Skin 1 had relatively more *Bacter-*

oidales and *Clostridiales* (c. 20–30%), while Skin 2 had greater abundances of the *Actinobacteridae* and *Bacillales* (c. 35–55%) (Fig. 2). These person-to-person differences in taxon abundance were also evident in the UniFrac analyses, as each sample clustered by host rather by temperature or length of storage (Fig. 1). Weighted pairwise UniFrac distances were significantly greater between the samples from the two individuals ($P < 0.001$) than between replicate subsamples stored at different temperatures ($P = 0.93$) or durations ($P = 0.53$). Similarly, we observed no significant effect on the phylogenetic diversity between replicate subsamples analyzed after 3 days of storage vs. 14 days of storage, irrespective of the storage temperature ($P > 0.05$ in all cases). The fact that the highly personalized nature of skin-associated bacterial communities (Gao *et al.*, 2007; Costello *et al.*, 2009; Grice *et al.*, 2009) was still apparent after 14 days at a range of temperatures, with storage conditions having relatively little impact on the community composition or diversity, has important implications for mass sampling efforts sponsored by various international human microbiome projects, which aim to relate microbial community structure and function to physiologic and pathophysiologic features in individuals with a range of




Fig. 2. The relative abundance of various bacterial taxa in fecal, skin and soil samples after 14 days of storage. Bars represent the mean abundance of each group at each temperature in descending order (e.g. 20, 4, -20 and -80 °C) with 1 SEM. Abundances of bacterial taxa were classified to family for the fecal samples and to order for the skin and soil samples using the RDP11 nomenclature. Differences in relative abundances due to storage temperature were assessed within individual samples using the Kruskal–Wallis test. * P -values ≤ 0.05 .

lifestyles in a variety geographic locations, some remote (Turnbaugh *et al.*, 2007).

Soil samples

The two soils harbored unique bacterial communities, with temperature and length of storage having little effect on the overall community composition (Figs 1 and 2, Table 1). Soils retained similar abundances of the six most numerous taxa across the range of storage temperatures tested, except for the *Burkholderiales*, which were marginally affected by temperature in Soil 1 ($P=0.05$, Fig. 2). Although each sample had similar abundances of most taxa, the two soil communities were clearly distinct regardless of the storage conditions ($P < 0.001$, Fig. 1 and Table 1). Analysis of UniFrac pairwise distances showed no significant effect of Day, Temperature or Day \times Temperature on the overall community composition for subsamples immediately frozen at -80 °C and those stored at 20 °C for 14 days ($P > 0.05$ in all cases). Likewise, phylogenetic diversity was unaffected by temperature or length of storage ($P > 0.05$ in all cases, Table 2).

Conclusion

We surveyed microbial communities from multiple environments under a broad range of storage conditions, and demonstrated that the bacterial community composition in the samples was largely unaffected by differences in short-term storage conditions. Although it is not currently possible to resolve changes in bacteria at the species or the strain level using pyrosequencing, given the limitations of read length and error rate (Kunin *et al.*, 2010), our results are consistent with other studies and indicate that the provenance of samples has a greater effect on the microbial community structure than do the conditions under which samples are stored before conducting DNA extractions. Critically, these differences persist both at a broad level (e.g. between soil and skin) and at the more subtle level of specific samples (e.g. different soils or skin from different people). Subsamples stored under different conditions did not have identical bacterial communities, perhaps due to insufficient sample homogenization or the inherent variability in DNA extractions and PCR amplification between subsamples. Importantly, these

Table 1. PERMANOVA results of sample, Day, Temperature and Day × Temperature effects on pairwise weighted and unweighted UniFrac distances

	d.f.	SS	MS	Pseudo- <i>F</i>	<i>P</i>
Unweighted UniFrac					
All samples					
Sample	5	26	5.1	25	0.001
Day	1	0.27	0.27	0.73	0.75
Temperature	3	0.64	0.21	0.56	1.0
Day × Temperature	3	0.66	0.22	0.57	1.0
Fecal					
Sample	1	3.3	3.3	19	0.001
Day	1	0.26	0.26	1.09	0.24
Temperature	3	0.58	0.19	0.79	0.83
Day × Temperature	3	0.57	0.19	0.77	0.86
Soil					
Sample	1	1.4	1.4	6.2	0.001
Day	1	0.27	0.27	1.11	0.15
Temperature	3	0.68	0.23	0.92	0.87
Day × Temperature	3	0.66	0.22	0.89	0.98
Skin					
Sample	1	1.0	1.0	4.8	0.001
Day	1	0.32	0.32	1.4	0.02
Temperature	3	0.67	0.22	0.97	0.61
Day × Temperature	3	0.72	0.24	1.05	0.22
Weighted UniFrac					
All samples					
Sample	5	14	2.8	301	0.001
Day	1	0.04	0.04	0.41	0.76
Temperature	3	0.05	0.02	0.15	1.0
Day × Temperature	3	0.04	0.01	0.12	1.0
Fecal					
Sample	1	0.83	0.83	101	0.001
Day	1	0.06	0.06	2.28	0.107
Temperature	3	0.07	0.02	0.93	0.475
Day × Temperature	3	0.03	0.01	0.38	0.894
Soil					
Sample	1	0.46	0.46	56	0.001
Day	1	0.04	0.04	2.07	0.11
Temperature	3	0.04	0.01	0.67	0.70
Day × Temperature	3	0.02	0.01	0.38	0.99
Skin					
Sample	1	0.87	0.87	79	0.001
Day	1	0.02	0.02	0.51	0.53
Temperature	3	0.03	0.01	0.28	0.93
Day × Temperature	3	0.05	0.02	0.53	0.73

Bold text indicates tests where $P < 0.05$.

d.f., degrees of freedom; SS, sum of squares; MS, mean squares.

other potential sources of variability were more important than the variability introduced by differences in storage temperature and duration between subsamples even after 14 days of storage at room temperature. Although specific taxa may change in relative abundance with different storage conditions, our data suggest that the types of samples in this study can be stored and shipped at room temperature without having a significant impact on the assessment of

Table 2. Faith's phylogenetic diversity (Faith's PD) analysis of samples

	Day 3	Day 14
Fecal 1	12 (0.9)	15 (0.7)
20 °C	9.2 (0.4)	15 (0.4)
4 °C	12 (0.1)	15 (0.8)
-20 °C	13 (0.3)	15 (0.3)
-80 °C	13 (0.3)	14 (0.4)
Fecal 2	7.7 (0.2)	7.9 (0.3)
20 °C	7.6 (0.3)	8.1 (0.4)
4 °C	7.7 (0.1)	7.6 (0.3)
-20 °C	7.6 (0.2)	8.2 (0.4)
-80 °C	7.8 (0.1)	7.8 (0.1)
Skin 1	13 (1.5)	13 (0.7)
20 °C	12 (0.2)	13 (0.7)
4 °C	12 (0.2)	14 (0.04)
-20 °C	13 (0.4)	13 (1.3)
-80 °C	16 (2.3)	12 (0.4)
Skin 2	8.2 (1.0)	7.6 (0.7)
20 °C	7.0 (0.1)	8.7 (0.8)
4 °C	8.1 (0.9)	7.0 (0.1)
-20 °C	7.8 (1.2)	7.8 (0.9)
-80 °C	9.7 (0.9)	6.9 (0.2)
Soil 1	38 (1.0)	39 (0.9)
20 °C	37 (0.9)	39 (1.2)
4 °C	36 (1.3)	39 (1.3)
-20 °C	39 (0.1)	40 (0.8)
-80 °C	40 (0.5)	40 (0.6)
Soil 2	43 (1.3)	40 (0.5)
20 °C	40 (0.6)	40 (0.1)
4 °C	43 (0.3)	39 (0.6)
-20 °C	44 (1.4)	39 (0.2)
-80 °C	45 (0.7)	40 (0.8)

Mean Faith's PD and SEM are shown for individual subsamples, and each temperature on both days. Temperature had a significant effect on Faith's PD for Fecal sample 1 on Day 3 and was significantly greater on Day 14 for Fecal 1 ($P < 0.05$ in both cases). The phylogenetic diversity of the remaining samples was unaffected by temperature or length of storage.

the overall community composition or the relative abundances of most major bacterial taxa.

Acknowledgements

We thank Donna Berg-Lyons for her help with the sample processing, Jill Manchester for her help with DNA sequencing, plus Micah Hamady and Elizabeth Costello for assistance with the bioinformatics analyses. We would also like to thank members of the Fierer lab group for help on previous drafts of this manuscript. This work was supported by grants from the National Science Foundation (EAR 0724960), the U.S. Department of Agriculture (2008-04346) (N.F.), the Howard Hughes Medical Institute (R.K.), the Bill and Melinda Gates Foundation, the Crohn's and Colitis Foundation of America and NIH (R01 HG004872) (R.K. and J.I.G.).

References

- Clarke KR & Warwick RM (2001) A further biodiversity index applicable to species lists: variation in taxonomic distinctness. *Mar Ecol-Prog Ser* **216**: 265–278.
- Cole JR, Chai B, Farris RJ, Wang Q, Kulam SA, McGarrell DM, Garrity GM & Tiedje JM (2005) The ribosomal database project (RDP-II): sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Res* **33**: D294–D296.
- Costello EK, Lauber CL, Hamady M, Knight R & Fierer N (2009) Bacterial variation in human body habitats across space and time. *Science* **326**: 1694–1697.
- DeSantis TZ, Hugenholtz P, Keller K, Brodie EL, Larsen N, Piceno YM, Phan R & Andersen GL (2006a) NAST: a multiple sequence alignment server for comparative analysis of 16S rRNA genes. *Nucleic Acids Res* **34**: W394–W399.
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P & Andersen GL (2006b) Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microb* **72**: 5069–5072.
- Dolfing J, Vos A, Bloem J, Ehlert PAI, Naumova NB & Kuikman PJ (2004) Microbial diversity in archived soils. *Science* **306**: 813a.
- Faith DP (1992) Conservation evaluation and phylogenetic diversity. *Biol Conserv* **61**: 1–10.
- Fierer N, Hamady M, Lauber CL & Knight R (2008) The influence of sex, handedness, and washing on the diversity of hand surface bacteria. *P Natl Acad Sci USA* **105**: 17994–17999.
- Gao Z, Tseng CH, Pei ZH & Blaser MJ (2007) Molecular analysis of human forearm superficial skin bacterial biota. *P Natl Acad Sci USA* **104**: 2927–2932.
- Grice EA, Kong HH, Conlan S *et al.* (2009) Topographical and temporal diversity of the human skin microbiome. *Science* **324**: 1190–1192.
- Hamady M & Knight R (2009) Microbial community profiling for human microbiome projects: tools, techniques, and challenges. *Genome Res* **19**: 1141–1152.
- Hamady M, Lozupone CA & Knight R (2009) Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *ISME J* **4**: 17–27.
- Hamady M, Walker JJ, Harris JK, Gold NJ & Knight R (2008) Error-correcting barcoded primers for pyrosequencing hundreds of samples in multiplex. *Nat Methods* **5**: 235–237.
- Klammer S, Mondini C & Insam H (2005) Microbial community fingerprints of composts stored under different conditions. *Ann Microbiol* **55**: 299–305.
- Kunin V, Engelbrekton A, Ochman H & Hugenholtz P (2010) Wrinkles in the rare biosphere: pyrosequencing errors can lead to artificial inflation of diversity estimates. *Environ Microbiol* **12**: 118–123.
- Lauber CL, Hamady M, Knight R & Fierer N (2009) Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microb* **75**: 5111–5120.
- Li WZ & Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.
- Lozupone C & Knight R (2005) UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microb* **71**: 8228–8235.
- Lozupone C, Hamady M & Knight R (2006) UniFrac - An online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinformatics* **7**.
- Lozupone CA & Knight R (2008) Species divergence and the measurement of microbial diversity. *FEMS Microbiol Rev* **32**: 557–578.
- Nechvatal JM, Ram JL, Basson MD, Namprachan P, Niec SR, Badsha KZ, Matherly LH, Majumdar APN & Kato I (2008) Fecal collection, ambient preservation, and DNA extraction for PCR amplification of bacterial and human markers from human feces. *J Microbiol Meth* **72**: 124–132.
- Ott SJ, Musfeldt M, Timmis KN, Hampe J, Wenderoth DF & Schreiber S (2004) *In vitro* alterations of intestinal bacterial microbiota in fecal samples during storage. *Diagn Microb Infect Dis* **50**: 237–245.
- Paulino LC, Tseng CH, Strober BE & Blaser MJ (2006) Molecular analysis of fungal microbiota in samples from healthy human skin and psoriatic lesions. *J Clin Microbiol* **44**: 2933–2941.
- Price MN, Dehal PS & Arkin AP (2009) FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol* **26**: 1641–1650.
- Rochelle PA, Cragg BA, Fry JC, Parkes RJ & Weightman AJ (1994) Effect of sample handling on estimation of bacterial diversity in marine sediments by 16S rRNA gene sequence analysis. *FEMS Microbiol Ecol* **15**: 215–225.
- Roesch LFW, Casella G, Simell O, Krischer J, Wasserfall CH, Schatz D, Atkinson MA, Neu J & Triplett EW (2009) Influence of fecal sample storage on bacterial community diversity. *Open Microbiol J* **3**: 40–46.
- Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R & Gordon JI (2007) The human microbiome project. *Nature* **449**: 804–810.
- Tzeneva VA, Salles JF, Naumova N, de Vos WA, Kuikman PJ, Dolfing J & Smidt H (2009) Effect of soil sample preservation, compared to the effect of other environmental variables, on bacterial and eukaryotic diversity. *Res Microbiol* **160**: 89–98.